

Visibility

Education and Research
from Envision

Depression and Vision Rehabilitation: Recognizing and Managing This Prevalent Co-morbidity..... Kara Crumbliss, OD

The primary roles of vision rehabilitation are to maximize visual function, facilitate adjustment to low vision and enhance quality of life. In order to do this effectively, providers must be knowledgeable about non-ocular conditions that commonly present in the low vision population. They must also adapt their examinations and encompass testing and treatment strategies in order to achieve successful rehabilitation and maintain quality of life for patients with compounding etiologies for their symptomatology. True clinical depression is a mood disorder in which feelings

of sadness, loss, anger or frustration interfere with everyday life for weeks or longer.¹ There are many contributing factors that may trigger onset of depression including significant change or stressful life events. Loss of vision may fall into either one of these categories. It seems intuitive that rates of depression would be higher amongst individuals dealing with vision loss than the general population. Indeed, this is well substantiated in published research. Older adults who are visually impaired report higher levels of depression than those who are not visually impaired,²⁻⁴ while

nearly half of patients with visual impairment who visit rehab centers report problems with emotional or psychological adjustment.⁵

Since both depression and vision loss may interfere with daily functioning and activities of daily living, treating vision loss without recognizing and treating depressive symptoms may not allow a patient to successfully meet their rehabilitation goals. This is supported by research showing that depression can prevent patients with low vision from achieving optimal rehabilitation outcomes.⁶⁻⁸ Additionally, depression has been reported

Feature Article

Practice Management Issues

Case Study

Professional Education

1 Depression and Vision Rehabilitation: Recognizing and Managing This Prevalent Co-morbidity

Kara Crumbliss, OD

6 Home Lighting Assessment and Modifications

Monica S. Perlmutter, MA, OTR/L, SCLV

10 Case Study: Doc, I Can See the Letters on Your Chart, But I Can't Read: Neuro-Rehabilitation Using the Visual Evoked Potential

Ronald Siwoff, OD, FAAO, DPL-ABO

14 Excellence in Education: Envision Conference 2012



Visibility is a quarterly publication of the Envision Foundation.

610 N. Main, Wichita, KS 67203
(316) 440-1600
www.ensonius.com

EDITORIAL STAFF

Michael Epp, MS, Director of Professional Education

Kelsey Rawson, Professional Education Associate

Shannon Riley, MA, Research and Analytics Associate

Kathi A. Buche, Graphic Design Manager

GUEST CONTRIBUTORS

Kara Crumbliss, OD

Monica S. Perlmutter, MA, OTR/L, SCLV

Ronald Siwoff, OD, FAAO, DPL-ABO

To submit an article or case study to be considered for publication in *Visibility*, please contact Michael Epp, Director of Professional Education, at (316) 440-1515 or michael.epp@ensonius.com.

The viewpoints expressed by the guest authors of *Visibility* do not necessarily reflect the viewpoints of Envision or its staff.

ABOUT ENVISION FOUNDATION

The mission of the Envision Foundation is to secure funding for the successful delivery of services offered by the Envision Vision Rehabilitation Center and the education programs of the Foundation. Envision Foundation focuses on fundraising to ensure that no patient is ever turned away—regardless of ability to pay; public education to help prevent blindness; and professional education to determine best practices in order to serve patients who are blind or low vision.

REQUEST COPIES OF VISIBILITY

If you would like to share *Visibility* with a colleague, please request a copy from Michael Epp, Director of Professional Education, at michael.epp@ensonius.com or call (316) 440-1515. *Visibility* is also available online at www.ensonius.com/Visibility.

Copyright © 2012 Envision Foundation. Individual articles are Copyright © 2012 of the indicated authors, printed with permission. All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or any information storage or retrieval system, without written permission of Envision Foundation.

Depression and Vision Rehabilitation... continued from front page

to be a more significant predictor for functional impairment than severity of vision loss.⁹ Individuals who report depression at the start of rehabilitation do not respond as well functionally to the goals of vision rehabilitation.⁸

Despite the preponderance of research documenting this relationship and the need for the co-management of these conditions for successful rehabilitation outcomes, only 11.8 percent of rehab centers in the United States have a psychologist on staff.⁵ This low number may be in response to financial practice management due to limited consensus and publication on reimbursement strategies for incorporating mental health professionals (MHPs), including social workers, psychologists and psychiatrists, into vision rehabilitation. Multidisciplinary management with outside referral of patients to an MHP is a viable option for patient management, yet only 31 percent of eye health professionals routinely screen for depression as part of patient management; conversely, 60 percent of rehabilitation workers do so.¹⁰ In this same study by Rees et al, knowledge seemed to be a key barrier, as 80 percent of eye health practitioners reported they would like to be more knowledgeable about signs and symptoms of depression.¹⁰ So it seems that while practitioners recognize they should screen for depression, few are doing so. Eye health professionals may be at a loss as to how and when to do this, and don't necessarily understand

the impact of overlooking this treatment on rehabilitation strategies.

Research has yet to firmly establish the change in outcomes for clinical depression and low vision treatments when combined,

example, a patient with relatively good acuity suffering from depression as a result of the stress of vision loss may have restored visual acuity and reading ability with a low powered magnifier in the



and to understand which, if either, should take priority. Yet, it is inferred that the symbiotic relationship would require treatment of both. Low vision rehabilitation and counseling may reduce risk of depression in individuals with vision impairment.¹¹ This antidepressant effect may benefit the practitioner and patient; however, the converse may also be true. Clinicians failing to identify and treat depression may result in poorer rehabilitation outcomes as depressive symptoms have been shown to correlate with perceived difficulty of activities of daily living and do not reliably correlate with levels of vision loss or worsening acuity.¹²⁻¹³ For

office, yet they may still claim they cannot read and it is too difficult as they perceive the difficulty of the task in the context of depression and not acuity. In this instance, the practitioner, failing to screen, recognize and refer for treatment of the depression, will be frustrated as the vision problem is not the only barrier to successful rehabilitation.

The American Academy of Ophthalmology, American Optometric Association, American Occupational Therapy Association and Association for Education and Rehabilitation of the Blind and Visually Impaired have endorsed the Optimum Low Vision Rehabilitation Service Delivery Model

which recognizes that support services, including counseling, community resources and support groups, form part of the optimum delivery model of vision rehabilitation services.¹⁴ Still, directives on best practice patterns were not



given in this model. Based on our knowledge that 80 percent of eye health practitioners would like to be more familiar with the signs and symptoms of depression, how do practitioners learn to recognize symptoms of depression?

Getting to know a local mental health practitioner who can provide counseling and depression treatment is imperative for successful practice. Generally, these social workers, psychologists or psychiatrists may not have firsthand experience working with low vision populations, just as eye health practitioners generally lack training and experience in management of depression. A sharing of information between these practitioners

can help the rehabilitation team members to understand the others' perspectives on working with individuals who have low vision. Screening tests for depression can be recommended by the counseling professional so that the eye health

care professional may use them and provide more appropriate and accurate referrals.


One suggested test is the Patient Health Questionnaire-9 (PHQ-9).¹⁵ This brief depression screening tool takes half the time of other depressive screening scales and can be particularly helpful in objective assessment of depressive symptoms. Better still, it has been validated in a small sample study of the low vision population, and was shown to screen for depression accurately and independently of vision loss.¹⁶ It is non-visual, can be administered by phone for those practices who wish to treat depression prior to or coincident to rehabilitation, and it does not

require a degree in psychology to score and interpret. This nine-item rating of state of depression (i.e., how someone is feeling now and in the past several days) on a four-point scale for each question, allows a five-point classification of depression, between mild and severe. Patients having moderate or worse levels of depression should be referred to an MHP qualified to evaluate and manage depression. Practitioners wishing to save examination time may also use the PHQ-2, which is only two questions; however, the PHQ-2, when positive, suggests administration of the PHQ-9 so practitioners must have both on hand in order to accurately complete the PHQ-2 for positive depressive response.¹⁷ For this reason, a personal preference is to habitually complete the PHQ-9 on patients. Anecdotally, since adding a psychologist to our staff, the simple addition of nine questions and an intake provided by an MHP to our patients, there has been an improvement in the rehabilitation outcomes and satisfaction of our patients.

A randomized clinical control to identify which tests, interventions, support groups or group, family or individual counseling, with or without adjunct antidepressant medical therapy, are appropriate for each patient, would help all practitioners incorporate an exact strategy for patient and practice management and is needed. Without such a study as yet published, practitioners can rely on shared experiences and suggest

best practice strategies just as this article does. Counseling services provided on-site may be billed for by the appropriately licensed and Medicare registered MHPs. In order to be most cost effective, a part-time or consultant MHP is, in many cases, a better business model in order to provide the service without additional expense to the practice.

Until the benefits of addressing psychosocial issues within vision rehabilitation plans of care are clearly established, it is unlikely that widespread incorporation of psychological support within

practices will occur. This review shows that recognizing and appropriately referring for depression is critical to the clinical practice of the vision rehabilitation field. Even non-rehabilitation eye care professionals who diagnose vision loss and its etiology should recognize depression and appropriately refer for treatment, in addition to referral for visual rehabilitation. This can have a positive impact on our patients' quality of life despite the presence of their eye diseases. 

References:

1. ADAM Medical Encyclopedia, 2012. <http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0001941/> Reviewed by: Zieve D, Merrill D. Review Date: 3/15/2011.
2. Evans JR, Fletcher AE, Wormald RP. Depression and anxiety in visually impaired older people. *Ophthalmology* 2007;114(2):283-8.
3. Bragg MW. Vision loss, depression and rehabilitation. *Int Congr Ser* 2005;1282:40-41.
4. Hayman KJ, Kerse NM, La Grow SJ, Woules T, Robertson MC, Campbell AJ. Depression in older people: Visual impairment and subjective ratings of health. *Optom Vis Sci* 2007;84(11):1024-30.
5. Owsley C, McGwin G Jr, Lee PP, Wasserman N, Searcey K. Characteristics of low-vision rehabilitation services in the United States. *Arch Ophthalmol* 2009;127(5):681-89.
6. Rovner BW, Casten RJ, Tasman WS. Effect of depression on vision function in age-related macular degeneration. *Arch Ophthalmol* 2002;120(8):1041-44.
7. Iliffe S, Kharicha K, Harari D, Swift C, Gillmann G, Stuck A. Self-reported visual function in healthy older people in Britain: An exploratory study of associations with age, sex, depression, education and income. *Fam Pract* 2005;22(6): 585-90.
8. Grant P, Seiple W, Szlyk JP. Effect of depression on actual and perceived effects of reading rehabilitation for people with central vision loss. *J Rehabil Res Dev* 2011;48(9):1101-8.
9. Banerjee A, Kumar S, Kulhara P, Gupta A. Prevalence of depression and its effect on disability in patients with age-related macular degeneration. *Indian J Ophthalmol* 2008; 56(6): 469-474.
10. Rees G, Fenwick EK, Keeffe JE, Mellor D, Lamoureux EL. Detection of depression in patients with low vision. *Optom Vis Sci* 2009;86(12):1328-36.
11. Horowitz A, Reinhardt JP, Boerner K. The effect of rehabilitation on depression among visually disabled older adults. *Aging Ment Health* 2005;9(6):563-70.
12. Rovner BW, Casten RJ, Hegel MT, Hauck WW, Tasman WS. Dissatisfaction with performance of valued activities predicts depression in age-related macular degeneration. *Int J Geriatr Psychiatry* 2007;22(8):789-93.
13. Casten RJ, Rovner BW. Vision Loss and Depression in the Elderly. *Psychiatric Times* 2006; 23(13) <http://www.psychiatrictimes.com/display/article/10168/>
14. Low Vision Rehabilitation Delivery Model Launched. ATeam representatives; Mogk L, Williams RT, Warren M, Deremeik J, Roberts D. August 10, 2007. <http://www.mdsupport.org/deliverymodel/deliverymodel.html>
15. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. Primary care evaluation of mental disorders. Patient Health Questionnaire. *JAMA* 1999; 282, 1737-44.
16. Lamoureux EL, Tee HW, Pesudovs K, Pallant JF, Keeffe JE, Rees G. Can clinicians use the PHQ-9 to assess depression in people with vision loss? *Optom and Vis Sci* 2009;86 (2), 139-145.
17. Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med Care* 2003 Nov;41(11):1284-92.

ACKNOWLEDGEMENTS

The author wishes to acknowledge the contributions of Patricia Grant, MS, PhD candidate, David Rakofsky, PsyD, and Alfred Rosenbloom, OD, FAAO, who co-presented portions of this material at Envision Conference 2011, September 23, 2011.

Dr. Crumbliss received her optometry degree from the Illinois College of Optometry in 2003 and completed a residency in Low Vision and Ocular Disease at the ICO at the Deicke Center for Visual Rehabilitation in 2004. Upon joining the faculty of the Illinois College of Optometry on a part-time basis in 2004, she also began practice as a Low Vision Consultant with The Chicago Lighthouse. Dr. Crumbliss is currently the Director of Clinical Services at The Chicago Lighthouse. She also works as the Coordinator of ICO's Low Vision and Ocular Disease Residency and as an Associate Clinical Professor in the Primary Care Service of the Illinois Eye Institute. Her research interests include the association of low vision with Charles Bonnet Syndrome, cognitive impairment and Alzheimer's.



Home Lighting Assessment and Modifications

..... *Monica S. Perlmutter, MA, OTR/L, SCLV*

Currently, approximately 2.4 million Americans have low vision and the prevalence of vision disabilities is expected to increase dramatically over the next 20 years.¹ Low vision is defined as a significant reduction in visual function that cannot be fully corrected with the use of eye glasses, contact lenses, or medical treatment, but is severe enough to interfere with daily activities. Low vision is associated with common age-related eye conditions seen in older adults, including macular degeneration and glaucoma. The growing population of older adults with vision loss is at risk for decline in ability to perform daily activities required to age in place.²

NORMAL AGE-RELATED CHANGES AFFECTING NEED FOR MORE LIGHT

As a person ages, the quality of their vision worsens due to reasons independent of age-related eye diseases. Several of these changes impact the amount of light that reaches the retina. Reduction in pupil size and the extent to which the pupil dilates decreases with age and allows less light to pass through. The aging lens and cornea causes glare due to light scattering. Slowing of dark adaptation can lead to night vision problems. By age 60, the lens transmits only 20 percent of light; if the person has cataracts, transmission is reduced to as low as 2 percent.³⁻⁵

RECOMMENDED LIGHTING LEVELS FOR OLDER ADULTS

The Illumination Engineering Society of North America (IESNA) recommends lighting levels for a range of domestic tasks performed in the home.⁶ For example, a healthy adult age 55 or older reading small print for prolonged periods requires a lighting level of approximately 50 footcandles or 540 lux to read small print for prolonged periods.⁶ General activities performed in the kitchen require 30 footcandles or 325 lux. Multiple studies show that home lighting levels are well below these recommended lighting levels.⁷⁻¹⁰

LIGHTING NEEDS OF PERSONS WITH AGE-RELATED OCULAR DISEASE

Individuals with ocular disease may require even higher levels of properly directed light. Study participants with macular degeneration experienced improvements in near visual acuity and contrast sensitivity with lighting levels increased from 300 to 3000 lux.¹¹ Eldred found that the majority of their sample with macular degeneration preferred illumination levels of 5920 lux or higher.¹² Lighting-related difficulties for individuals with glaucoma include problems with glare intolerance, dark/light adaptation and difficulty with low-lit environments.¹³

Cataracts may cause poor vision outside with glare due to sunlight, and during night driving due to headlights and lamps.¹⁴ Persons with glaucoma and cataracts may benefit from additional lighting that is properly directed (i.e. below eye level, directly over reading material) to avoid problems with glare. Unfortunately, environmental lighting in the community cannot be easily modified.

BENEFITS OF OPTIMAL LIGHTING

Several studies have shown that optimal lighting leads to increased visual function.¹⁵⁻¹⁷ One study showed that increasing the amount of light had a significant effect on sentence reading acuity, reading rate and critical print size for persons with macular degeneration.¹⁵ Brunstrom and colleagues¹⁶ examined the effect of improvement in lighting on activities of daily living, quality of life and well-being and found that higher lighting levels led to



ENVISION

improvement in pouring and slicing bread and increased quality of life. In another study, participants were able to see and recognize more objects in a simulated living room with higher lighting levels.¹⁷ Despite the known benefits of lighting, older adults may not fully appreciate the role of lighting and may rate their current home lighting as adequate.¹⁸ Furthermore, improving lighting, contrast and safety may be viewed as less important than learning daily living skill strategies.¹⁹ Older adults and persons with ocular disease would benefit from learning how efficient lighting can promote the greatest use of their residual vision and how inadequate home lighting may lead to loss of activities and pose significant safety concerns.

HOME LIGHTING ASSESSMENT AND OCCUPATIONAL THERAPY

Home lighting should be included in occupational therapy assessment and intervention for a number of key reasons. The literature has established that the aging eye allows reduced amounts of light to reach the retina and that home lighting is well below recommended levels. It is also well documented that many persons with ocular disease require significant increases in lighting and some ocular diseases cause light sensitivity. Lindner and colleagues¹⁰ found that more than 70 percent (54/76) of study participants used ceiling lights versus task lights as primary lighting sources for



Home Lighting Intervention: Under counter task lighting (top); Ambient lighting – bedroom before and after (middle row); Lighting enhancement–bathroom (bottom left), hallway (bottom middle) and closet (bottom right)

reading, and only 40 percent used additional task lighting. Given that older adults spend an average of 12 to 16 hours, or 80 percent of their day, in their own homes,²⁰ it is clear that home lighting is a critical area for occupational therapists to address.

Lighting preferences are very individual, thus, assessment approaches should include objective and subjective assessment of visual comfort to determine optimal illumination.^{15,17,21} Several standard measures include items related

to home lighting. The Housing Enabler includes 180 items related to the physical home environment – two of which relate to outdoor lighting and illumination of walkways.²² The Vision Version of the In Home Occupational Performance Evaluation (I-HOPE) includes assessment of quantity, position and direction of lighting, glare, color, contrast and light changes with day, night or season, however it is only available in French.²³ The Craig Hospital Inventory of Environmental Factors (CHIEF) is a

25-item questionnaire used to identify environmental barriers to participation. The CHIEF was created for large scale research purposes for use in the general population, but is not standardized for use with individuals with low vision.²⁴ Lighting is assessed in broad terms and is included along with other environmental factors such as noise and crowds. The Home Occupation Environment Assessment (HOEA) is a measure of home safety, and includes items related to physical accessibility, sanitation, proper food storage, general safety and lighting levels at the point of task.²⁵ Lighting levels are measured with a light meter at the location where the person reads, pays bills, prepares meals or performs other tasks of daily living.

EVIDENCE GUIDING COMPONENTS OF HOME LIGHTING ASSESSMENT


Existing measures do not fully assess the key elements that should be included in a home environment lighting assessment for older adults with low vision; the literature provides evidence that highlights components that should be incorporated. Certainly, the most obvious element to include is an objective measure of illumination. Recommended levels of illumination from IESNA⁶ and other literature^{7,8} can serve as a guide regarding optimal levels for specific diagnoses, such as macular degeneration.^{11,12} Use of a light meter in the home environment will allow the low vision specialist

to gather objective data about illumination levels. Glare intolerance and sources of direct and indirect glare should be included in a home lighting assessment. Glare reduces visual comfort and can be controlled with use of filters, window treatments, and covering reflective table and counter tops. Assessment of seating arrangement and positioning of reading material must be considered.²⁶ Qualitative aspects and comfort within the lighting environment are also important to assess. Four of the 18 study participants in Eldred's study¹² experienced discomfort and eye strain with use of high illumination levels for prolonged periods of time. Bowers, Meek and Stewart found that the ideal level of lighting determined by objective means was higher than levels determined by subjective preference, most likely because of greater visual comfort.¹⁵ Eye strain, satisfaction with lighting environment, ability to tolerate prolonged periods of reading/near-task activity and overall comfort level should also be addressed when evaluating a home lighting environment. A new measure, the Home Environment Lighting Assessment (HELA), is currently being developed by the author and addresses each of these components. The HELA will be available within the coming year; those who are interested can contact the author.²⁷

LIGHTING INTERVENTION STRATEGIES

A variety of texts and resources provide general guidelines for lighting intervention strategies which are very helpful, including *Lighting the Way: A Key to Independence and Low Vision Rehabilitation: Practical Guide for Occupational Therapists*.^{28,29} These resources encourage us to consider overall room and task lighting, to increase the wattage, move the light closer, and reduce glare. For reading, the light should be positioned over the shoulder; for writing, position the light opposite the person's dominant hand. Lights should be dimmed to watch TV or use the computer; paths should be lit from the bed to bathroom door with nightlights. When proposing lighting modifications, clients should be given the opportunity to compare the effect of optimal and sub-optimal lighting.³⁰ In addition, clients should be allowed to compare their reading performance at different lighting levels because there is variability among individuals.²¹ These techniques are logical, but are largely untested, with the exception of increasing light levels as previously described. Potential barriers to making lighting modifications include the resident's resistance to change, preference for a certain type of fixture or décor, limited financial resources to purchase lighting and lack of transportation to shop for fixtures.²⁹ Common concerns related to use of daylight as a means to boost lighting levels also exist. Older adults may prefer

to keep their shades and blinds closed due to security reasons; blinds and shades may not work properly and others may experience discomfort from the glare or draft.¹⁸

In summary, it is important to assess home lighting in a comprehensive, objective fashion so that we can develop lighting intervention plans that are tailored to the individual.^{12,31} Further research examining the efficacy of lighting modifications is warranted. 

Monica S. Perlmutter, MA, OTR/L, SCLV, is an instructor in the Occupational Therapy Program at Washington University, and is the lead OT for the program's Community Practice Low Vision Program. Her research studies have focused on the development of a measurement model to assess the impact of vision loss on older adults and examination of the affect of glaucoma on daily activities and quality of life. A new line of research aims to develop a standard method of assessing home lighting and to examine the efficacy of approaches to lighting modifications.



References:

- Congdon N, O'Colmain B, Klaver CC, Klein R, Muñoz B, Friedman DS, Kempen J, Taylor HR, Mitchell P; Eye Diseases Prevalence Research Group. Causes and prevalence of visual impairment among adults in the United States. *Arch Ophthalmol*. 2004; 122: 477-85.
- Berger S, Porell, F. The association between low vision and function. *J Aging Health* 2008; 20(5): 504-525.
- National Eye Institute. (n.d.). *What you should know about low vision*. Retrieved December 26, 2011, from <http://www.nei.nih.gov/health/lowvision/>.
- Jackson GR, Owsley C, McGwin G. Aging and dark adaptation. *Vision Res* 1999; 39:3975-3982.
- Sadun AA, Libondi T. Transmission of light through cataracts. *Am J Ophthalmol* 1990; 10(6): 710-712.
- Illumination Engineering Society of North America Guidelines (n.d.). Retrieved February 29, 2012 from: <http://www.ibacos.com/high-performance-lighting-guide/iesna-guidelines>.
- Charness N, Dijkstra K. Age, luminance, and print legibility in homes, offices, and public places. *Hum Factors* 1999; 41(2):173-193.
- Cullinan TR, Silver JH, Gould ES, Irvine D. Visual disability and home lighting. *Lancet* 1979; 1(8117): 642-644.
- Levitt JG. Lighting for the elderly: An optician's view. Paper read at the Symposium on Light for Partial Sight. Held by the Illuminating Engineering Society and The Partially Sighted Society in London, April, 1978.
- Lindner H, Rinnert T, Behrens-Baumann W. Illumination conditions of visually impaired people under private domestic conditions. *Klin Monatsbl Augenh* 2001; 218(12): 744-781.
- Haymes SA, Lee J. Effects of task lighting on visual function in age-related macular degeneration. *Ophthalmic & physiological optics*. *Ophthalmic Physiol Opt* 2006; 26(2):169-79.
- Eldred KB. Optimal illumination for reading in patients with age-related maculopathy. *Optom Vis Sci* 1992; 69(1):46-50.
- Nelson P, Aspinall P, Papasouliotis O, Worton B, O'Brien C. Quality of life in glaucoma and its relationship with visual function. *J Glaucoma* 2003; 12(2):139-150.
- National Eye Institute (n.d.) *Facts about cataracts*. Retrieved December 23, 2011, from http://www.nei.nih.gov/health/ataract/ataract_facts.asp.
- Bowers AR, Meek C, Stewart N. Illumination and reading performance in age-related macular degeneration. *Clin Exp Optom* 2001; 84(3):139-147.
- Brunnström G, Sörensen S, Alsterstad K, Sjöstrand J. Quality of light and quality of life—The effect of lighting adaptation among people with low vision. *Ophthalm Physiol Opt* 2004; 24(4):274-280.
- Cornelissen W, Bootsma A, Kooijman AC. Object perception by visually impaired people at different light levels. *Vision Res* 1995; 35(1):161-168.
- Bakker R, Iofel Y, Lachs, MS. Lighting levels in the dwellings of homebound older adults. *J Housing for the Elderly* 2004; 18(2):17-27.
- Schuchard RA, Naseer S, de Castro K. Characteristics of AMD patients with low vision receiving visual rehabilitation. *J Rehabil Res Dev* 1999; 36(4):294-302.
- Horgas AL, Wilms HU, Baltes MM. Daily life in very old age: Everyday activities as expression of successful living. *Gerontologist* 1998; 38(5): 556-568.
- Fosse P, Valberg A. Lighting needs and lighting comfort during reading with age-related macular degeneration. *J Vis Impair Blind* 2004; 98: 389-409.
- Slaug, B (2001). Enabler FAQ. Retrieved from <http://www.enabler.nu/>.
- Carigman M, Rousseau J, Gresset J, Couturier JA. Content validity of a home-based person-environment interaction assessment tool for visually impaired adults. *J Rehabil ResDev* 2008; 45(7):1037-1052.
- Whiteneck GG, Harrison-Felix CL, Mellick DC, Brooks CA, Charlifue SB, Gerhart KA. Quantifying environmental factors: A measure of physical, attitudinal, service, productivity, and policy barriers. *Arch Phys Med Rehabil* 2004; 85(8):1324-1335.
- Baum CM, Edwards DF. Guide for the Home Occupational-Environmental Assessment. 1988; St. Louis, MO: Washington University Program in Occupational Therapy.
- Markowitz M. Occupational Therapy Interventions in Low Vision Rehabilitation. *Can J Ophthalmol* 2006; 41(3):340-347.
- Perlmutter MS. The Home Environment Lighting Assessment (Unpublished doctoral thesis), 2012; Washington University, St. Louis, MO.
- Figueiro MG. *Lighting the way: A key to independence*. 2001; Troy, NY: Rensselaer Polytechnic Institute.
- Scheiman M, Sheiman M, Whittaker S. *Low vision rehabilitation: A practical guide for occupational therapists*. 2007; Thorofare, NJ: Slack Incorporated.
- Slay DH. Home-based environmental lighting assessments for people who are visually impaired: developing techniques and tools. *J Vis Impair Blind* 2002; 109-114.
- LaGrow SJ. Assessing optimal illumination for visual response accuracy in visually impaired adults. *J Vis Impair Blind* 1986; 8: 888-895.

Case Study: Doc, I Can See the Letters on Your Chart, But I Can't Read: Neuro-Rehabilitation Using the Visual Evoked Potential.....

Ronald Siwoff, OD, FAAO, DPL-ABO

The Visually Evoked Potential (VEP) is an objective measure of the functioning of the human visual system. It measures the strength of response to the signal (amplitude) as well as the time it takes for the signal to arrive at the primary visual cortex.¹⁻² The VEP uses a computer to select electrical signals from the brain that originate from the eye. VEP has been used in the past, primarily with infants and young children, to determine if they have sight.³⁻⁵ Now we find that the technology can reveal secrets of how the human visual system processes information.

The patient ("A.A.") presented with the complaint that she could not read. Distance VA was tested

with The Original Distance Chart for the Partially Sighted, arranged by William Feinbloom, OD, PhD. Near VA was established with The Lighthouse Near Acuity Test (Second Edition), Modified ETDRS with Sloan Letters. The patient had sufficient acuities for reading small print. Yet, when she was given text of various sizes, she could not read text at any size, large or small. This finding prompted a VEP evaluation.

Gold cup electrodes (Grass Model F-E56 H Astro-Med Inc., West Warwick, R.I.) were applied to the scalp with EEG paste. The electrodes were placed 4 cm above the inion on midline and 4 cm above the brow on midline. The ground electrode was placed midway

between the other electrodes. The impedance of the electrodes, measured prior to testing, was between 6 and 10 K-ohms.

A Diopsys Infant™ System (Diopsys Inc., Pine Brook, New Jersey, USA) was used with a checkerboard reversal pattern at 85% contrast viewed at one meter. First the check size with the biggest amplitude is selected.

128 X 128	4 cycles/degree
64 X 64	2 cycles/degree
32 X 32	1 cycles/degree
16 X 16	.533 cycles/degree

The contrast was set to Michelson 85%. The mean luminance was 66.25 cd/m². The checkerboard reversed at two per second. The testing time was 20 seconds.


Testing was performed monocularly with best corrected spectacle correction. The four check sizes were analyzed to determine which produced the largest amplitude. This check size was used four more times with ophthalmic trial frame prisms. The protocol used an 8 prism diopters, base up, base right, base down and base left. The N50, N75 and P100 were all identified. The amplitude was measured by subtracting the N50 from the P100. The orientation of the prism that resulted in the largest amplitude was selected. If two orientations

resulted in large amplitudes, the prescribed orientation was placed between both original orientations. For example, if the signals peaked in the Base-In and Base-Up orientations, the prescribed prism for the patient was both BU and BI. If three orientations resulted in large amplitudes, the orientation was placed 180° away from the smallest amplitude. If there was no improvement, no prism correction was prescribed. In the case of "A.A.", 8 prism diopters was the size of prisms in both eyes. In other patients, other sizes of prisms are used, depending on a formula used with acuities. Although, in the case of "A.A.", the prism is the same in both eyes, i.e., same size and direction, this is not always the case; both eyes do not necessarily take the same prism.

The signal must arrive at the primary visual cortex by 100 msec. If responses to signals are delayed, i.e., there is a latency in response, empty information arrives at Broca's area and, therefore, cannot be converted to language, which is necessary for reading and speech. One way to understand this is to think of a train leaving the station every 100 msec. If the train arrives late, empty boxcars will be sent to the next stop.

RESULTS

When tested with only the conventional correction, the patient, "A.A.", demonstrated significant latencies in right and left eyes. Prisms decreased latencies in both eyes. The right eye went from 110.0 to 101.6 msec, and the left eye



	T=20-P=32-OS-85%-C-12/29/11 6:57:35 PM	T=20-P=32-OS-85%-C-12/29/11 6:46:59 PM
Left Cursor	51.66 ms	53.33 ms
Right Cursor	185.0 ms	113.3 ms
Delta	53.33 ms	60.00 ms
Eye	OS	OS
CheckSize	32	32
Test Time	20	20
Contrast	85	85
Pattern	Checkerboard	Checkerboard
Binasal	N	N
Correction	Y	N
Comments	8 base down	

Copyright (c) 2009 Diopsys, Inc., All Rights Reserved



went from 113.3 to 105.0 msec. Once prism was added to the correct refraction, "A.A." could read normal print easily.

IMPRESSIONS AND CONCLUSION

Our ongoing research with a large sample of patients has demonstrated that ophthalmic prisms can change VEP amplitudes and latencies and improve visual acuity and function over

and above conventional spectacle prescriptions. This method has applications for anyone with mild to severe alexia caused by cortical damage or optic nerve disease.* Our outcomes suggest that when responses to signals are delayed, Broca's area cannot process the information needed to convert the signals into words and sentences, and therefore reading effectively is not possible.



PATIENT OCULAR HISTORY

Patient ID: "A.A." 91 YO WF
Occupation: Retired executive secretary
CC: Cannot read
DHx: Age-related macular degeneration, cataracts removed in both eyes
MHx: Hypertension, mild dementia, colon cancer
Allergies: Scallops, aspirin

VA Without Rx: OD 10/20-3 1.25M-1
 OS 10/20-1 1.25M-1

VA With Rx: OD 10/20-3 .8M
 OS 10/20-3 .8M

Refraction: OD +1.25 = -2.50 X 90 10/20
 OS +0.50 = -2.00 X 105 10/20
 ADD +6.00 OD .63M
 OS .8M

IOP: OD 15 mm Hg
 OS 15 mm Hg

Retina Evaluation:

OD – The cup/disc ratio was .3. Margins were clear. Peripapillary atrophy was noted. RPE atrophy was noted in the macula. The vein/artery ratio was 2/1. Atherosclerosis was noted. No holes, tears, atrophy or degeneration was noted in the peripheral retina to the ora serrata.
 OS – No significant difference from OD observed.

Anterior Segment Evaluation:

OD – The cornea was clear. Arcus senilis was noted 360° around the limbus. No staining with fluorescein was noted. No cell or flare was noted. A well-positioned posterior intraocular lens was noted. The peripheral lens capsule was opacified.
 OS – No significant difference from OD observed.


Visual Evoked Potential Evaluation Without Prisms: Abnormal

P-100 OD 110.0
 OS 113.3

Visual Evoked Potential With Prisms:

P-100 OD 101.6
 OS 105.0

Final Rx With Prisms: OD +1.25 = -2.50 X 90, 8 PD, DN
 OS +0.50 = -2.00 X 105, 8 PD, DN
 ADD: +6.00



	T=20-P=8-OD-85%-C-12/29/11 6:44:47 PM	T=20-P=8-OD-85%-C-12/29/11 6:41:12 PM
Left Cursor	50.00 ms	40.00 ms
	-0.6 uV	-0.3 uV
Right Cursor	101.6 ms	110.0 ms
	5.17 uV	4.63 uV
Delta	51.66 ms	70.00 ms
	2.78 uV	5.02 uV
Eye	OD	OD
CheckSize	8	8
Test Time	20	20
Contrast	85	85
Pattern	Checkerboard	Checkerboard
Binasal	N	N
Correction	Y	N
Comments	8 base down	

Copyright (c) 2009 Diopsys, Inc., All Rights Reserved



Dr. Siwoff is a member of the National Physician's Advisory Board, from which he received the 2004 Physician of the Year award. Other awards for his service to the blind and visually impaired community include the Outstanding Scientific Achievement award from the New Jersey Society of Optometric Physicians and the New Jersey Department of Human Services award for Outstanding Accomplishments and Contributions to the Blind and Visually Impaired Community. Prior to his present position as Director of the Siwoff Low Vision Center, Dr. Siwoff was Clinical Assistant Professor of Ophthalmology at the University of Medicine and Dentistry of New Jersey, where he directed the Low Vision Clinic. He later became Director of the Gerald E. Fonda Low Vision Center at Saint Barnabas Hospital in Livingston, New Jersey. Dr. Siwoff holds several patents for technologies to improve the vision of the visually impaired. He has conducted joint research with the NIH and Harvard Medical School.



References:

1. Odom JV, et al. Visual evoked potentials standard. *Doc Ophthalmol* 2004; 108:115-123.
2. Sokol S. Visually evoked potentials: theory, techniques and clinical applications. *Surv Ophthalmol* 1976 Jul-Aug;21(1):18-44.
3. Taylor MJ, McCulloch DL. Visual evoked potentials in infants and children. *J Clin Neurophysiol* 1992; 9(3):357-72.
4. Simon JW, et al. A new visual evoked potential system for vision screening in infants and young children. *J AAPOS* 2004; 8:549-554
5. Wygnanski-Jaffe T, Pantoni CM, Buncic JR, Westall CA. Paradoxical robust visual evoked potentials in young patients with cortical blindness. *Doc Ophthalmol* 2009; 119(2):101-7. Epub 2009 Jun 23.

*The author is preparing a full-length research paper on these results of evolved technical application of VEP.

Meet Me in St. Louis!

ENVISION CONFERENCE 2012

On behalf of the Envision Conference staff, we invite you to take part in the seventh annual Envision Conference, to be held September 12-15, 2012 in historic St. Louis, Missouri. The Envision Conference is a multi-disciplinary low vision rehabilitation and research conference dedicated to improving the quality of low vision care through professional collaboration, advocacy, research and education.



PROMOTE YOUR COMPANY. By sponsoring, exhibiting or advertising, you have the rare opportunity to share your message or products and services with hundreds of low vision professionals involved in the clinical and research arenas.

Exhibit

Advance Price: \$850 (paid on or before July 6, 2012)

Regular Price: \$950 (paid after July 6, 2012)

Premium Price: \$1,400 (only three available, paid during registration)

EXPAND YOUR KNOWLEDGE. Register today! Registration is now open online at www.envisionconference.org. Register by June 29 and save \$100 off regular registration.

Registration Fees

Regular: \$475 by 6/29; \$575 after 6/29; \$625 onsite

Student: \$225 by 6/29; \$250 after 6/29; \$260 onsite

Guest: \$100

One Day Pass: \$275

Pre-conference Workshops: \$100/each

Sponsor

Visionary Sponsor – Exclusive: \$10,000

Welcome Reception Sponsor: \$8,000

Buffet Lunch Sponsor: \$5,000

Speaker Reception Sponsor – Exclusive: \$5,000

Exhibit Hall Game Sponsor – Exclusive: \$4,000

Tote Bag Sponsor – Exclusive: \$4,000

Conference Program Sponsor – Exclusive: \$3,000

Conference Badge Lanyard Sponsor – Exclusive: \$2,500

Internet Cafe Sponsor: \$2,500

Advertise

Full Page Ad: \$500

Half Page Ad: \$300

Flyer in Attendee Bags: \$500

Take One Display: \$200

For more information on these great marketing opportunities, visit the Envision Conference website.

PLAN YOUR TRIP. St. Louis offers more free major attractions for the entire family than anyplace outside the nation's capital. In St. Louis, the Zoo (one of the finest in the world), Art Museum, Science Center, History Museum, Anheuser-Busch Brewery tours, Grant's Farm, Botanical Gardens and a host of other stops do not charge admission. The city is filled with trendy shops, charming neighborhoods, exceptional arts, as well as many historical, cultural and entertaining family attractions with top-notch restaurants. St. Louis has it all.

The Hilton St. Louis at the Ball Park is offering discounted room rates to Envision Conference 2012 attendees for \$139 per night. Reservations can be made by calling 1-877-845-7354. When calling, make sure to say you are with the group Envision Conference. An online reservation link is also available at www.envisionconference.org on the Hotel & Travel page. Please book your room early; this rate is only valid until August 17, 2012 or when the Envision block of rooms sells out, whichever comes first.





COMING SOON



Envision is pleased to announce the launch of Envision University.

The mission of Envision University is to provide multi-disciplinary continuing education and research opportunities for low vision rehabilitation professionals, establishing best practices to ensure continued research and clinical care for individuals who are blind or visually impaired. Our promise is to collaborate with vision rehabilitation and research professionals to provide relevant multi-disciplinary continuing education and research opportunities that address practice gaps in current standards of care and research.

Envision University will be comprised of live continuing education events, the annual Envision Conference, *Visibility* and expanded online education and research opportunities. Stay tuned for more information!



Envision Professional Education Calendar

April 12, 2012

Low Vision Grand Rounds – Current Medical Treatment and Vision Rehabilitation Options for Wet Macular Degeneration. Wichita, KS. CE – ACCME, AOTA, COPE

June 15, 2012

Neurological Vision Loss. Wichita, KS. CE – AOTA, COPE

July 12, 2012

Low Vision Grand Rounds – What’s New and Exciting in Corneal Surgery. Wichita, KS. CE – ACCME, AOTA, COPE

September 12-15, 2012

Envision Conference 2012. Hilton St. Louis at the Ballpark, St. Louis, MO. CE – ACCME, ACVREP, AOTA, CRCC, COPE

October 11, 2012

Low Vision Grand Rounds – Research on Contact Lenses. Wichita, KS. CE – ACCME, AOTA, COPE

November 9, 2012

Assistive Technology. Wichita, KS. CE – ACVREP, AOTA, CRCC, COPE

For more information, visit the Education and Resources page at www.envisionus.com.